



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## GEOMETRY.

80. Proposed by J. C. GREGG, Superintendent of Schools, Brazil, Ind.

One circle touches another internally, and a third circle whose radius is a mean proportional between their radii, passes through the point of contact. Prove that the other intersections of the third circle with the first two are in a line parallel to the common tangent of the first two. [From *Phillips and Fisher's Geometry*.]

I. Solution by G. B. M. ZERR, A. M., Ph. D., President and Professor of Mathematics in Russell College, Lebanon, Va., and the PROPOSER.


In the two triangles  $CDB$  and  $CDA$ , the angle  $CDG$  is common; also  $DB : DC = DC : DA$ .

$\therefore$  The two angles are similar and  $\angle BCD = \angle CAD$ ,  $\angle CBD = \angle ACD$ . But  $\angle DCB = \angle DEG$ , both being measured by arc  $DK$ .

$\therefore \angle DEG = \angle CAD$ .

Since  $DE$  is perpendicular to  $CA$ ,  $EG$  must be perpendicular to  $CA$ .

$\therefore HD$  and  $EG$  are parallel.



II. Solution by HENRY HEATON, M. Sc., Atlantic, Iowa, and J. SCHEFFER, A. M., Hagerstown, Md.

Taking the common tangent of the first two circles as the axis of  $x$  and the common diameter as the axis of  $y$  and supposing the tangent of the third circle, through the origin to make the angle  $\alpha$  with the axis of  $x$ , the equations of the circles are,

$$x^2 + y^2 - 2ay = 0,$$

$$x^2 + y^2 - 2by = 0, \text{ and}$$

$$x^2 + y^2 - 2c(x\sin\alpha + y\cos\alpha) = 0.$$

If the first and third circles intersect,

$$y = 0 \text{ or } \frac{2ac^2\sin^2\alpha}{a^2 + 2accos\alpha + c^2}.$$

If the second and third circles intersect,

$$y = 0 \text{ or } \frac{2bc^2\sin^2\alpha}{b^2 + 2bccos\alpha + c^2}.$$

If the line through the second intersections is parallel to the axis of  $x$  the two values of  $y$  are equal.

$\therefore ab^2 + ac^2 = a^2b + bc^2$ . Whence  $c^2 = ab$ .

If  $c = \sqrt{ab}$ , the distance  $y$  of the intersections from the common tangent is in both cases

$$\frac{2absin^2\alpha}{a + \sqrt{ab}\cos\alpha + ab}.$$

Q. E. D.

Also solved by COOPER D. SCHMITT and CHARLES C. CROSS.